

WEYL ALMOST PERIODIC SELECTIONS OF MULTIVALUED MAPS

L.I. DANILOV

Let (U, ρ) be a complete metric space and let a point $x_0 \in U$ be fixed. For $p \geq 1$, we denote by $M_p(\mathbb{R}, U)$ the set of (strongly) measurable functions $f : \mathbb{R} \rightarrow U$ such that

$$\sup_{\xi \in \mathbb{R}} \int_{\xi}^{\xi+1} \rho^p(f(t), x_0) dt < +\infty.$$

On $M_p(\mathbb{R}, U)$ we introduce the metrics

$$D_{p,l}^{(\rho)}(f, g) = \left(\sup_{\xi \in \mathbb{R}} \frac{1}{l} \int_{\xi}^{\xi+l} \rho^p(f(t), g(t)) dt \right)^{1/p}, \quad l > 0,$$

and the semimetric

$$D_p^{(\rho)}(f, g) = \lim_{l \rightarrow +\infty} D_{p,l}^{(\rho)}(f, g), \quad f, g \in M_p(\mathbb{R}, U).$$

For a measurable set $T \subseteq \mathbb{R}$ (in the sense of Lebesgue), we use the notation

$$\varkappa_W(T) = \lim_{l \rightarrow +\infty} \sup_{\xi \in \mathbb{R}} \frac{1}{l} \text{meas} [\xi, \xi + l] \cap T,$$

where *meas* is Lebesgue measure on $\mathbb{R}$. Let $M_p^*(\mathbb{R}, U)$ be the set of functions $f \in M_p(\mathbb{R}, U)$ such that for any $\varepsilon > 0$ there is a number $\delta > 0$ such that for all measurable sets $T \subseteq \mathbb{R}$ with $\varkappa_W(T) < \delta$ the inequality

$$\lim_{l \rightarrow +\infty} \sup_{\xi \in \mathbb{R}} \frac{1}{l} \int_{[\xi, \xi+l] \cap T} \rho^p(f(t), x_0) dt < \varepsilon$$

holds.

Let $\varepsilon > 0$. For a function $f \in M_p(\mathbb{R}, U)$, a number $\tau \in \mathbb{R}$ is called $(\varepsilon, D_{p,l}^{(\rho)})$ -almost period if $D_{p,l}^{(\rho)}(f(\cdot), f(\cdot + \tau)) < \varepsilon$, and is called $(\varepsilon, D_p^{(\rho)})$ -almost period if $D_p^{(\rho)}(f(\cdot), f(\cdot + \tau)) < \varepsilon$. A set $T \subseteq \mathbb{R}$ is *relatively dense* if there exists a number $a > 0$ such that $[\xi, \xi + a] \cap T \neq \emptyset$ for all $\xi \in \mathbb{R}$. A function $f \in M_p(\mathbb{R}, U)$, $p \geq 1$, belongs to the space $W_p(\mathbb{R}, U)$ of *Weyl almost periodic (a.p.)* functions of *order* p if for any $\varepsilon > 0$ there is a number $l = l(\varepsilon, f) > 0$ such that the set of $(\varepsilon, D_{p,l}^{(\rho)})$ -almost periods of f is relatively dense.

On the space U , we also define the metric $\rho'(x, y) = \min\{1, \rho(x, y)\}$, $x, y \in U$. Let $W(\mathbb{R}, U) \doteq W_1(\mathbb{R}, (U, \rho'))$ be the space of *Weyl a.p.* functions $f : \mathbb{R} \rightarrow U$ (of order 1)

taking values in the metric space (U, ρ') . We have

$$W_p(\mathbb{R}, U) \subseteq W_1(\mathbb{R}, U) \subseteq W(\mathbb{R}, U), \quad p \geq 1.$$

A sequence $\{\tau_j\}_{j \in \mathbb{N}} \subset \mathbb{R}$ is said to be *f-returning* for a function $f \in W(\mathbb{R}, U)$ if $D_1^{(\rho')}(f(\cdot), f(\cdot + \tau_j)) \rightarrow 0$ as $j \rightarrow +\infty$. For a function $f \in W(\mathbb{R}, U)$, by $\text{Mod } f$ we denote the module (additive group) of all numbers $\lambda \in \mathbb{R}$ such that $e^{i\lambda\tau_j} \rightarrow 1$ as $j \rightarrow +\infty$ (where $i^2 = -1$) for all *f-returning* sequences $\{\tau_j\}$. If $U = (\mathcal{H}, \|\cdot\|)$ is a Banach space and $f \in W_1(\mathbb{R}, \mathcal{H})$, then $\text{Mod } f$ coincides with the frequency module of the function f .

Let $(\text{cl}_b U, \text{dist}_\rho)$ be the metric space of non-empty closed bounded sets of the space (U, ρ) with the Hausdorff metric dist_ρ ; $(\text{cl } U, \text{dist}_{\rho'})$ is the metric space of non-empty closed sets of the space (U, ρ) with the Hausdorff metric $\text{dist}_{\rho'}$ corresponding to the metric ρ' . The spaces $W_p(\mathbb{R}, \text{cl}_b U)$, $p \geq 1$, and $W(\mathbb{R}, \text{cl}_b U)$ of *Weyl a.p. multivalued maps* $F : \mathbb{R} \rightarrow \text{cl}_b U$ are defined as the spaces of Weyl a.p. functions taking values in the metric space $(\text{cl}_b U, \text{dist}_\rho)$. Let $W(\mathbb{R}, \text{cl } U) \doteq W_1(\mathbb{R}, (\text{cl } U, \text{dist}_{\rho'}))$; $W(\mathbb{R}, \text{cl}_b U) \subseteq W(\mathbb{R}, \text{cl } U)$.

Theorem 1 (see [1]). *Let (U, ρ) be a complete metric space, let $g \in W(\mathbb{R}, U)$, and let $F \in W(\mathbb{R}, \text{cl } U)$. Then for any increasing function $\eta : [0, +\infty) \rightarrow [0, +\infty)$, for which $\eta(0) = 0$ and $\eta(t) > 0$ for all $t > 0$, there is a function $f \in W(\mathbb{R}, U)$ such that $\text{Mod } f \subseteq \text{Mod } g + \text{Mod } F$, $f(t) \in F(t)$ a.e., and*

$$\rho(f(t), g(t)) \leq \rho(g(t), F(t)) + \eta(\rho(g(t), F(t))) \text{ a.e.}$$

If $F \in W_p(\mathbb{R}, \text{cl}_b U)$ for some $p \geq 1$, then the function f belongs to the space $W_p(\mathbb{R}, U)$.

For a complete separable metric space (U, ρ) we denote by $(\mathcal{M}(U), d)$ the (complete separable) metric space of Borel probability measures $\mu[\cdot]$ defined on the σ -algebra of Borel subsets of the metric space (U, ρ) , with the Lévy – Prokhorov metric d ; $W_1(\mathbb{R}, \mathcal{M}(U))$ is the space of *Weyl a.p. measure-valued functions* $\mathbb{R} \ni t \rightarrow \mu[\cdot; t] \in \mathcal{M}(U)$ (of order 1) taking values in the metric space $(\mathcal{M}(U), d)$. A measure-valued function $\mathbb{R} \ni t \rightarrow \mu[\cdot; t] \in \mathcal{M}(U)$ belongs to the space $W_1(\mathbb{R}, \mathcal{M}(U))$ if and only if

$$\int_U \mathcal{F}(x) \mu[dx; \cdot] \in W_1(\mathbb{R}, \mathbb{R})$$

for all continuous bounded functions $\mathcal{F} \in C_b(U, \mathbb{R})$ (see [2]). Moreover,

$$\text{Mod } \mu[\cdot; \cdot] = \sum_{\mathcal{F} \in C_b(U, \mathbb{R})} \text{Mod } \int_U \mathcal{F}(x) \mu[dx; \cdot].$$

For $\mu \in \mathcal{M}(U)$, $x \in U$, and $\delta \in (0, 1)$ we shall use the notation

$$r_\delta(x, \mu) = \inf \{r > 0 : \mu[U_r(x)] > \delta\},$$

where $U_r(x) = \{y \in U : \rho(x, y) < r\}$.

Theorem 2. *Let (U, ρ) be a complete separable metric space, let $g \in W(\mathbb{R}, U)$, and let $\mu[\cdot; \cdot] \in W_1(\mathbb{R}, \mathcal{M}(U))$. Then for any $\delta \in (0, 1)$ there exists a function $f_\delta \in W(\mathbb{R}, U)$ such that $\text{Mod } f_\delta(\cdot) \subseteq \text{Mod } g(\cdot) + \text{Mod } \mu[\cdot; \cdot]$, $f_\delta(t) \in \text{supp } \mu[\cdot; t]$ a.e., and*

$$\rho(f_\delta(t), g(t)) < r_\delta(g(t), \mu[\cdot; t]) + \delta \text{ a.e.}$$

Let $\mathcal{S}_{\text{rd}}$ be the collection of relatively dense subsets $T \subseteq \mathbb{R}$. We shall denote by $\mathcal{P}_p^{(\rho)}(\varepsilon; f)$ the set of $(\varepsilon, D_p^{(\rho)})$ -almost periods of a function $f \in M_p(\mathbb{R}, U)$. Let $\widetilde{W}_p(\mathbb{R}, U)$ be the set of functions $f \in M_p(\mathbb{R}, U)$ satisfying the following two conditions:

- 1) $\mathcal{P}_p^{(\rho)}(\varepsilon; f) \in \mathcal{S}_{\text{rd}}$ for all $\varepsilon > 0$;
- 2) for any $\varepsilon, \delta > 0$ there is a compact set $K_{\varepsilon, \delta} \subseteq U$ such that $\varkappa_W(\{t \in \mathbb{R} : \rho(f(t), K_{\varepsilon, \delta}) \geq \varepsilon\}) < \delta$. (In the case $U = (\mathbb{R}^n, |\cdot|)$, the condition 2 is satisfied for all functions $f \in M_1(\mathbb{R}, \mathbb{R}^n)$.)

Let $\widetilde{W}(\mathbb{R}, U) \doteq \widetilde{W}_1(\mathbb{R}, (U, \rho'))$. We have

$$W_p(\mathbb{R}, U) = W(\mathbb{R}, U) \cap M_p^*(\mathbb{R}, U) \subseteq \widetilde{W}(\mathbb{R}, U) \cap M_p^*(\mathbb{R}, U) \subseteq \widetilde{W}_p(\mathbb{R}, U).$$

If $f \in \widetilde{W}(\mathbb{R}, U) \cap M_p^*(\mathbb{R}, U)$, then for any $\varepsilon' > 0$ there exists a number $\varepsilon > 0$ such that $\mathcal{P}_p^{(\rho)}(\varepsilon'; f) \supseteq \mathcal{P}_1^{(\rho')}(\varepsilon; f)$.

Theorem 3. *Let (U, ρ) be a complete metric space, let $g \in \widetilde{W}(\mathbb{R}, U)$, and let $F \in \widetilde{W}_1(\mathbb{R}, (\text{cl } U, \text{dist}_{\rho'}))$. Suppose that*

$$\mathcal{T}(\varepsilon) \doteq \mathcal{P}_1^{(\rho')}(\varepsilon; g) \cap \mathcal{P}_1^{(\text{dist}_{\rho'})}(\varepsilon; F) \in \mathcal{S}_{\text{rd}}$$

for all $\varepsilon > 0$. Then for any $\delta > 0$ and any function $h \in W_1(\mathbb{R}, \mathbb{R})$ with a dense (countable) frequency module $\text{Mod } h$ (for this function $\mathcal{T}(\varepsilon) \cap \mathcal{P}_1^{(\rho_{\mathbb{R}})}(\varepsilon; h) \in \mathcal{S}_{\text{rd}}$ for all $\varepsilon > 0$) there exists a function $f_\delta \in \widetilde{W}(\mathbb{R}, U)$ such that $f_\delta(t) \in F(t)$ a.e.,

$$\rho(f_\delta(t), g(t)) < \rho(g(t), F(t)) + \delta \text{ a.e.},$$

and for any $\varepsilon' > 0$ there is a number $\varepsilon > 0$ such that

$$\mathcal{P}_1^{(\rho')}(\varepsilon'; f_\delta) \supseteq \mathcal{T}(\varepsilon) \cap \mathcal{P}_1^{(\rho_{\mathbb{R}})}(\varepsilon; h).$$

Furthermore, if $F \in M_p^(\mathbb{R}, \text{cl}_b U)$ for some $p \geq 1$, then $f_\delta \in \widetilde{W}(\mathbb{R}, U) \cap M_p^*(\mathbb{R}, U) \subseteq \widetilde{W}_p(\mathbb{R}, U)$.*

REFERENCES

- [1] L. I. Danilov, *On Weyl almost periodic selections of multivalued maps*, J. Math. Anal. Appl. **316** (2006), 110–127.
- [2] L. I. Danilov, *On Weyl almost periodic measure-valued functions*, Izv. Inst. Mat. i Inform. Udmurt. Univ., vyp. 1 (31), Izhevsk, 2005, pp. 79–98 (in Russian).

PHYSICAL-TECHNICAL INSTITUTE, URAL BRANCH OF THE RUSSIAN ACADEMY OF SCIENCES, KIROV STREET 132, IZHEVSK 426000, RUSSIA

E-mail address: danilov@otf.pti.udm.ru